

***Misdiagnosed conditions prior to Morquio A diagnosis<sup>30</sup>***

| <b><i>Misdiagnosis</i></b>     | <b><i>N (%)</i></b> |
|--------------------------------|---------------------|
| Spondyloepiphiseal dysplasias  | 5 (28)              |
| Gibbus/kyphosis                | 5 (28)              |
| Scoliosis                      | 3 (17)              |
| Craniosynostosis               | 2 (11)              |
| Advanced bone/dental age       | 2 (11)              |
| Overgrowth                     | 2 (11)              |
| Genu valgum                    | 2 (11)              |
| Corneal opacity                | 2 (11)              |
| Cardiac conduction abnormality | 2 (11)              |
| Legg-Calvé-Perthes disease     | 1 (6)               |
| Leri-Weill syndrome            | 1 (6)               |
| Marfan syndrome                | 1 (6)               |

|                             |       |
|-----------------------------|-------|
| Sotos syndrome              | 1 (6) |
| Pseudoachondroplasia        | 1 (6) |
| Truncal hypotonia           | 1 (6) |
| Hypertelorism               | 1 (6) |
| Splenomegaly                | 1 (6) |
| Growth hormone deficiency   | 1 (6) |
| Torticollis                 | 1 (6) |
| Sleep obstruction (snoring) | 1 (6) |
| Cervical lymphadenopathy    | 1 (6) |
| Apnoea/hypopnoea            | 1 (6) |
| Rheumatoid arthritis        | 1 (6) |
| Harrison sulcus             | 1 (6) |
| Lumbar lordosis             | 1 (6) |
| Reversed madelung deformity | 1 (6) |

Adapted from Bhattacharaya, *Orphanet J Rare Dis*, 2014.