Ongoing multisystemic assessments in patients with Morquio ${\bf A}^3$

Asssessments	Frequency
Musculoskeletal	
Standardised upper extremity function test, radiographs	At diagnosis/baseline, annually
Respiratory	
FVC, MVV, respiratory rate, oxygen saturation, overnight sleep study	At diagnosis/baseline, annually
Neurological	
Neurological exam	 At diagnosis/baseline, every visit (minimum, every 6 months)
Plain radiograph MRI scan	• At diagnosis, every 1 to 3 years
CT scan	At diagnosis, annually
C1 Scall	As clinically indicated
Cardiovascular	
Electrocardiogram	 At diagnosis, every 1 to 3 years, as clinically indicated
Echocardiogram	 At diagnosis, every 2 to 3 years, as clinically indicated
Heart rate	At diagnosis, annually
Ophthalmological	
Refractive error and intraocular pressure	At diagnosis, as clinically indicated
Audiological	
Multimodal hearing assessment	At diagnosis, annually
Abdominal	
Assessments of gastrointestinal problems	As clinically indicated
Dental	
Evaluation of oral health	At diagnosis, annually
Endurance	
6MWT	\bullet At diagnosis, annually, before and regularly after initiation of ERT
QoL	
Reproducible, age-appropriate QoL questionnaires (e.g. EQ-SD-SL)	At diagnosis, annually
	,,

Adapted from Hendriksz, Am I Med Genet Part A,2014.

Abbreviations: 6MWT, 6-minute walk test; CT, computed tomography; ERT, enzyme replacement therapy; FVC, forced vital capacity; MRI, magnetic resonance imaging; MVV, maximum voluntary ventilation; QoL, quality of life.